

Particle-like solutions to topologically massive gravity

Gérard Clément*

Laboratoire de Gravitation et Cosmologie Relativistes

Université Pierre et Marie Curie, CNRS/URA769

Tour 22-12, Boîte 142

4, place Jussieu, 75252 Paris cedex 05, France

March 25, 1994

Abstract

The solution of topologically massive gravity with cosmological constant is reduced, for space-times with two commuting Killing vectors, to a special-relativistic dynamical problem. This approach is applied to the construction of a class of exact sourceless, horizonless solutions asymptotic to the BTZ extreme black holes.

*E-mail: GECL@CCR.JUSSIEU.FR.

Three-dimensional Einstein gravity with a negative cosmological constant has recently been shown by Bañados, Teitelboim and Zanelli (BTZ) [1] to admit black-hole solutions which are in some respects quite similar to the Kerr black holes of four dimensional general relativity. There is however an important difference between the two theories: the field equations of three-dimensional cosmological gravity imply that the curvature is constant outside sources, so that the theory is dynamically trivial. To remedy this defect, Deser, Jackiw and Templeton [2] have proposed a theory of three-dimensional gravity with a Chern-Simons topological self-coupling — topologically massive gravity (TMG) — which admits massive excitations. The field equations of TMG with cosmological constant are

$$G^\mu{}_\nu + \frac{1}{\mu} C^\mu{}_\nu = \Lambda \delta^\mu{}_\nu, \quad (1)$$

where $G^\mu{}_\nu \equiv R^\mu{}_\nu - \frac{1}{2} R \delta^\mu{}_\nu$ is the Einstein tensor,

$$C^\mu{}_\nu \equiv \varepsilon^{\mu\alpha\beta} D_\alpha (R_{\beta\nu} - \frac{1}{4} g_{\beta\nu} R) \quad (2)$$

is the Cotton tensor (with $\varepsilon^{\mu\alpha\beta}$ the antisymmetrical tensor), μ is the topological mass constant (we shall assume without loss of generality $\mu > 0$) and Λ is the cosmological constant. As remarked by Kaloper [3], these equations are trivially solved by the BTZ black-hole metric, which has a covariantly constant Ricci tensor, leading to a vanishing Cotton tensor¹.

Is there a non-trivial TMG analogue of the four-dimensional Kerr solution? Let us first make more precise what we mean by a Kerr analogue. We would like such a solution 1) to have an event horizon, and 2) for $\Lambda \leq 0$, to be asymptotically particle-like, which we define as asymptotic to the conical metric of [4] for $\Lambda = 0$, or asymptotic to the BTZ metric for $\Lambda < 0$. The black-hole solution of TMG with cosmological constant recently given by Nutku [8] satisfies only the first of these criteria; this solution (eq. (18) below), which is a generalization of the $\Lambda = 0$ Vuorio solution [9], does have an event horizon for a certain range of parameter values, but is not asymptotically particle-like. On the other hand, two kinds of asymptotically particle-like wormhole

¹Note that this holds only because the BTZ black hole is everywhere regular. The well-known point-particle solution to three-dimensional gravity [4] solves TMG only when the mass m and spin σ of the source are constrained by $m + \mu\sigma = 0$ [5, 6, 7].

solutions have been found in TMG with vanishing cosmological constant [10], but these do not have horizons. In this Letter we shall construct a family of asymptotically particle-like solutions to TMG with negative cosmological constant. These new solutions turn out not to have a regular horizon, and so do not satisfy our first criterion. However a subclass of these space-times are geodesically complete, and asymptotic to the BTZ extreme black holes, which we feel qualifies them as candidates for particle-like solutions of TMG. After all, what is the necessity of a horizon if there is no singularity to hide?

We first present a new, powerful approach to reduce the solution of TMG with two commuting Killing vectors K_1, K_2 to a dynamical problem in an abstract three-dimensional Minkowski space. This approach is based on the observation [10, 11] that the $SL(2, R)$ group of transformations in the plane (K_1, K_2) is locally isomorphic to the Lorentz group $SO(2, 1)$, which suggests the parametrization of the “physical” three-dimensional metric

$$ds^2 = \lambda_{ab}(\rho) dx^a dx^b + \zeta^{-2}(\rho) R^{-2}(\rho) d\rho^2, \quad (3)$$

where λ is the 2×2 matrix

$$\lambda = \begin{pmatrix} T + X & Y \\ Y & T - X \end{pmatrix}, \quad (4)$$

$\det \lambda = R^2 \equiv \mathbf{X}^2$ is the Minkowski pseudo-norm of the vector $\mathbf{X} = (T, X, Y)$,

$$\mathbf{X}^2 \equiv T^2 - X^2 - Y^2 \quad (5)$$

(special linear transformations of λ correspond to Lorentz rotations of $\mathbf{X}$), and the function $\zeta(\rho)$ allows for arbitrary reparametrizations of the coordinate ρ . If the vector $\mathbf{X}$ is space-like ($R^2 < 0$), the physical metric (3) has the Minkowskian signature, and is stationary rotationally symmetric if the orbits of the space-like Killing vector are closed.

The action for TMG with cosmological constant Λ is

$$I = I_E + I_{CS}, \quad (6)$$

where

$$I_E = -m \int d^3x \sqrt{|g|} (g^{\mu\nu} R_{\mu\nu} + 2\Lambda) \quad (7)$$

is the action for cosmological gravity ($\kappa = 1/2m$ is the Einstein gravitational constant), and

$$I_{CS} = -\frac{m}{2\mu} \int d^3x \varepsilon^{\lambda\mu\nu} \Gamma_{\lambda\sigma}^\rho (\partial_\mu \Gamma_{\rho\nu}^\sigma + \frac{2}{3} \Gamma_{\mu\tau}^\sigma \Gamma_{\nu\rho}^\tau) \quad (8)$$

is the Chern-Simons term. The action (6) reduces with the parametrization (3) to

$$I = \int d^2x \int d\rho L, \quad (9)$$

where the generalized Lagrangian [12]

$$L = -\frac{m}{2} \zeta \dot{\mathbf{X}}^2 - 2m \zeta^{-1} \Lambda + \frac{1}{2\mu m} \zeta^2 (\mathbf{X}, \dot{\mathbf{X}}, \ddot{\mathbf{X}}) \quad (10)$$

depends on the coordinates X^i , velocities $\dot{X}^i = dX^i/d\rho$, accelerations $\ddot{X}^i$ and on the dynamical variable ζ , which acts as a Lagrange multiplier (in (10), $(\mathbf{X}, \dot{\mathbf{X}}, \ddot{\mathbf{X}})$ is the mixed product $\varepsilon_{ijk} X^i \dot{X}^j \ddot{X}^k$). Varying this Lagrangian with respect to the coordinates X^i leads to the equations of motion

$$\ddot{\mathbf{X}} = -\frac{1}{2\mu} (3 \dot{\mathbf{X}} \wedge \ddot{\mathbf{X}} + 2 \mathbf{X} \wedge \dot{\ddot{\mathbf{X}}}). \quad (11)$$

These dynamical equations do not involve the cosmological constant, which occurs only in the constraint

$$H \equiv -\frac{1}{2} m \dot{\mathbf{X}}^2 + \frac{m}{\mu} (\mathbf{X}, \dot{\mathbf{X}}, \ddot{\mathbf{X}}) = -2m \Lambda \quad (12)$$

obtained by varying the Lagrangian (10) with respect to ζ (in (11) and (12), we have set the arbitrary function ζ equal to 1 after variation). Equations (11) and (12) are equivalent to the field equations (1) in the case of a metric depending only on one variable.

The action (9) is by construction invariant under pseudo-rotations and under translations in ‘time’ ρ . The invariance under rotations leads to the conservation of generalized angular momentum

$$\mathbf{J} = m \mathbf{X} \wedge \dot{\mathbf{X}} + \frac{m}{2\mu} [2 \mathbf{X} \wedge (\mathbf{X} \wedge \ddot{\mathbf{X}}) - \dot{\mathbf{X}} \wedge (\mathbf{X} \wedge \dot{\mathbf{X}})], \quad (13)$$

while the constant of the motion associated with translations of ρ is the generalized hamiltonian H occurring in the left hand-side of eq. (12), which is of course the hamiltonian constraint for TMG.

Solutions of Einstein cosmological gravity ($\mu \rightarrow \infty$) depending on only one variable are geodesics of our abstract Minkowski space,

$$\mathbf{X} = \boldsymbol{\alpha} \rho + \boldsymbol{\beta}, \quad (14)$$

with $\boldsymbol{\alpha}^2 = 4\Lambda$ from the Hamiltonian constraint; obviously these also solve the TMG equations (11), (12) for finite μ . In this Letter we make the following ansatz for solutions to TMG:

$$\mathbf{X} = \boldsymbol{\alpha} \rho + \boldsymbol{\beta} + \boldsymbol{\gamma} \rho^p \quad (15)$$

($\rho > 0, p$ real, $p \neq 0, 1$). Then eqs. (11) imply that

$$\boldsymbol{\alpha} \wedge \boldsymbol{\gamma} = -\frac{2\mu}{2p-1} \boldsymbol{\gamma} \quad (16)$$

($p \neq 1/2$), and either 1) $p = 2$, or 2) $\boldsymbol{\beta} \wedge \boldsymbol{\gamma} = 0$. Taking into account the relations $\boldsymbol{\alpha}^2 = -4\mu^2/(2p-1)^2$, $\boldsymbol{\alpha} \cdot \boldsymbol{\gamma} = \boldsymbol{\gamma}^2 = 0$ which follow from (16), the Hamiltonian constraint (12) then leads to the relation

$$\frac{\mu^2}{(2p-1)^2} - \frac{p(p-1)}{2p-1} (\boldsymbol{\beta} \cdot \boldsymbol{\gamma}) \rho^{p-2} = -\Lambda. \quad (17)$$

In the first case ($p = 2$), the solution depends a priori on five parameters, three for the vectors $\boldsymbol{\alpha}$, $\boldsymbol{\gamma}$ related by (16), two for the vector $\boldsymbol{\beta}$ restricted by (17). However, as in the case of Einstein cosmological gravity [11], stationary rotationally symmetric solutions really depend on only two parameters, as three of the parameters may be chosen at will by suitable metric reparametrizations involving time dilations, uniform frame rotations, and translations of ρ (under which the ansatz (15) with $p = 2$ is form-invariant). The parametrization

$$\boldsymbol{\alpha} = (-1, 1, -\frac{2\mu}{3}), \quad \boldsymbol{\beta} = (a, a, \frac{3b}{\mu}), \quad \boldsymbol{\gamma} = \frac{\mu^2 + 9\Lambda}{12a} (1, -1, 0)$$

leads to the metric

$$\begin{aligned} ds^2 = & 2a dt^2 - 2 \left(\frac{2\mu}{3} \rho - \frac{3b}{\mu} \right) dt d\theta + \left(\frac{\mu^2 + 9\Lambda}{6a} \rho^2 - 2\rho \right) d\theta^2 \\ & - \frac{d\rho^2}{\frac{\mu^2 - 27\Lambda}{9} \rho^2 + 4(a-b)\rho + \frac{9b^2}{\mu^2}}, \end{aligned} \quad (18)$$

which reduces to Nutku's solution [8] if one defines $M = 12(b-a)$, $J = 6b$, $r^2 = 2\rho$ (and takes care that the cosmological constant in [8] is $-\Lambda$). This generalized Vuorio solution has horizons for certain ranges of parameter values, but is obviously not asymptotically particle-like.

The second possibility ($\beta \wedge \gamma = 0$) leads to $\beta \cdot \gamma = 0$ (γ is null), so that the constraint (17) can be satisfied only if the cosmological constant is negative. Putting $\Lambda = -l^{-2}$ ($l > 0$), this constraint selects the two possible values of the exponent p ,

$$p_{\pm} = \frac{1}{2} (1 \pm \mu l). \quad (19)$$

The constant angular momentum (13) is then related to the null vector β by

$$\mathbf{J} = \frac{2m}{\mu l^2} (1 \pm \mu l) \beta. \quad (20)$$

Now the general solution depends a priori on four arbitrary parameters (three for the vectors α , γ related by (16), one for the constant c determining $\beta = c^{-1}\gamma$), but the ansatz (15) is not form-invariant under translations of ρ , so that stationary rotationally symmetric solutions depend again on two parameters. We choose the parametrization

$$\alpha = l^{-2} (1 - l^2, 1 + l^2, 0), \quad \beta = -\frac{M}{4} (1 + l^2, 1 - l^2, \mp 2l), \quad \gamma = c\beta,$$

such that the metric

$$\begin{aligned} ds^2 = & \left(2l^{-2} \rho - \frac{m_{\pm}(\rho)}{2} \right) dt^2 \pm l m_{\pm}(\rho) d\theta dt \\ & - \left(2\rho + l^2 \frac{m_{\pm}(\rho)}{2} \right) d\theta^2 - \frac{l^2 d\rho^2}{4\rho^2} \end{aligned} \quad (21)$$

with

$$m_{\pm}(\rho) = M (1 + c \rho^{p_{\pm}}) \quad (22)$$

reduces to the BTZ extreme black hole [1] for $c = 0$, $M > 0$.

The solutions (21) are asymptotically particle-like if $p_{\pm} < 0$, which occurs only for the exponent p_- in the case $\mu l > 1$. The corresponding solutions are asymptotic to the BTZ extreme black-hole metric for $M > 0$, or to the corresponding extreme point-particle metric for $M < 0$. The metric (21) is apparently singular at $\rho = 0$, which is at infinite proper distance. To check whether $\rho = 0$ is also at infinite geodesic distance, we write the first integral of the (physical) geodesic equation

$$R^{-2} \left(\frac{d\rho}{d\tau} \right)^2 + \Pi^T \lambda^{-1} \Pi = \varepsilon, \quad (23)$$

where Π is a constant column matrix, τ is an affine parameter, and $\varepsilon = +1, 0$ or -1 . With the parametrization (5), eq. (23) simplifies to

$$\left(\frac{d\rho}{d\tau} \right)^2 + \mathbf{P} \cdot \mathbf{X} - \varepsilon R^2 = 0, \quad (24)$$

where $\mathbf{P}$ is a constant null vector. In the present case this equation reads

$$\left(\frac{d\rho}{d\tau} \right)^2 + \boldsymbol{\alpha} \cdot \mathbf{P} \rho + \boldsymbol{\beta} \cdot \mathbf{P} (1 + c \rho^{p_{\pm}}) + \varepsilon \frac{4\rho^2}{l^2} = 0. \quad (25)$$

For $p_- < 0$, the effective potential near $\rho = 0$ is dominated by the term $\boldsymbol{\beta} \cdot \mathbf{P} c \rho^p$, which is attractive or repulsive depending on the sign of cM . For $cM > 0$, the constant $c\boldsymbol{\beta} \cdot \mathbf{P}$ is generically negative, so that $\rho = 0$ is at finite geodesic distance; unless $c = 0$ (in which case $\rho = 0$ is the horizon of the BTZ extreme black hole), the geodesics cannot for p_- real be extended to negative ρ , so the space-time of metric (21) is singular² on the circle $\rho = 0$. On the other hand, for $cM < 0$, $c\boldsymbol{\beta} \cdot \mathbf{P} \geq 0$ and ρ can attain the value 0 only if the null vectors $\mathbf{P}$ and $\boldsymbol{\beta}$ are collinear (implying also $\boldsymbol{\alpha} \cdot \mathbf{P} = 0$)

²This space-time is singular even though the curvature invariants $R = 6/l^2$, $R^{\mu\nu} R_{\mu\nu} = 12/l^4$ are regular (another instance in TMG is the exact $f = 0$ solution of [10], eq. (27) below). Indeed, the singularity is due to the term $\boldsymbol{\gamma} \rho^p$ in (15), which does not contribute to the curvature invariants because the vector $\boldsymbol{\gamma}$ is null and orthogonal to $\boldsymbol{\alpha}$ and $\boldsymbol{\beta}$.

and $\varepsilon = -1$; the time-like circle $\rho = 0$ is at infinite affine distance on these space-like geodesics, so that the geometry (21) is regular (as in the case of the extreme point- particle solution of Einstein cosmological gravity [11]).

For completeness, we also briefly discuss the case $p_{\pm} > 0$, which occurs for p_+ if $\mu l > 1$, or for both exponents $p_{\pm}$ if $\mu l < 1$. Now the effective potential near $\rho = 0$ is dominated by the constant $\boldsymbol{\beta} \cdot \boldsymbol{P}$. We do not repeat the discussion, which follows closely that made in the case $p_- < 0$, leading to the conclusion that the geometry (21) is singular for $M > 0$, and regular for $M < 0$.

The solution (21) is such that the vector $\boldsymbol{X}(\rho)$ evolves in the plane $(\boldsymbol{\alpha}, \boldsymbol{\beta})$ orthogonal to the constant vector $\boldsymbol{J}$. One may inquire whether there are more general planar solutions

$$\boldsymbol{X} = \boldsymbol{\alpha} F(\rho) + \boldsymbol{\beta} G(\rho) \quad (26)$$

to cosmological TMG. We have found that this planar family of solutions includes, besides the trivial solution (14) (Λ real) and the solution (21) ($\Lambda < 0$), only one other solution, the exact $f = 0$ solution of [10] ($\Lambda = 0$),

$$\boldsymbol{X} = \boldsymbol{\alpha} + \boldsymbol{\beta} (c \rho + e^{-\nu \rho}), \quad (27)$$

which has properties somewhat similar to those of solution (21), and is particle-like for $c = 0$.

We have reduced the solution of TMG with two commuting Killing vectors to a dynamical problem in an abstract three-dimensional Minkowski space. A polynomial ansatz leads to two classes of solutions. The Vuorio-Nutku solutions of the first class ($p = 2$) are not asymptotically particle-like. The second class ($p = \frac{1}{2}(1 \pm \mu l)$) includes a subclass of regular solutions asymptotic to the BTZ extreme black holes — sourceless, horizonless particle-like solutions of TMG with negative cosmological constant.

Acknowledgement

I thank S. Deser for stimulating discussions and helpful suggestions.

References

- [1] Bañados M, Teitelboim C and Zanelli J 1992 *Phys. Lett.* **69** 1849
Bañados M, Henneaux M, Teitelboim C and Zanelli J 1993 *Phys. Rev. D* **48** 1506
- [2] Deser S, Jackiw R and Templeton S 1982 *Phys. Rev. Lett.* **48** 975; 1982 *Ann. Phys., NY* **140** 372
- [3] Kaloper N 1993 *Phys. Rev. D* **48** 2598
- [4] Deser S, Jackiw R and 't Hooft G 1984 *Ann. Phys., NY* **152** 220
- [5] Deser S 1990 *Phys. Rev. Lett.* **64** 611
- [6] Ortiz M E 1990 *Class. Quant. Grav.* **7** L13
- [7] Clément G 1990 *Class. Quant. Grav.* **7** L193
- [8] Nutku Y 1993 *Class. Quant. Grav.* **10** 2657
- [9] Vuorio I 1985 *Phys. Lett.* **163B** 91
Percacci R, Sodano P and Vuorio I 1987 *Ann. Phys., NY* **176** 344
- [10] Clément G 1992 *Class. Quant. Grav.* **9** 2615
- [11] Clément G 1994 *Phys. Rev. D*, in press
- [12] Goldstein H 1980 *Classical Mechanics* (Reading, Mass.: Addison-Wesley), p.66